



## NEVADA CITY SCHOOL OF THE ARTS

### Procedures for Informing Parents about Diabetes

#### **Type 2 Diabetes: Information for Parents of 7th Grade Students**

Type 2 diabetes is the most common form of diabetes. Millions of Americans have been diagnosed with type 2 diabetes, and many more are unaware they are at high risk. In type 2 diabetes, either the body does not produce enough insulin or the cells ignore the insulin. Insulin is necessary for the body to be able to use glucose for energy. When you eat food, the body breaks down all of the sugars and starches into glucose, which is the basic fuel for the cells in the body. Insulin takes the sugar from the blood into the cells. When glucose builds up in the blood instead of going into cells, it can lead to health complications.

#### **Risk Factors Associated with Type 2 Diabetes**

It is recommended that students displaying or possibly experiencing the risk factors and warning signs associated with type 2 diabetes be screened (tested) for the disease.

#### **Risk Factors**

Researchers do not completely understand why some people develop type 2 diabetes and others do not; however, the following risk factors are associated with an increased risk of type 2 diabetes in children:

**Being overweight.** The single greatest risk factor for type 2 diabetes in children is excess weight. In the U.S., almost one out of every five children is overweight. The chances are more than double that an overweight child will develop diabetes.

**Family history of diabetes.** Many affected children and youth have at least one parent with diabetes or have a significant family history of the disease.

**Inactivity.** Being inactive further reduces the body's ability to respond to insulin.

**Specific racial/ethnic groups.** Native Americans, African Americans, Hispanics/Latinos, or Asian/Pacific Islanders are more prone than other ethnic groups to develop type 2 diabetes.

**Puberty.** Young people in puberty are more likely to develop type 2 diabetes than younger children, probably because of normal rises in hormone levels that can cause insulin resistance during this stage of rapid growth and physical development.

#### **Warning Signs and Symptoms Associated with Type 2 Diabetes**

Warning signs and symptoms of type 2 diabetes in children develop slowly, and initially there may be no symptoms. However, not everyone with insulin resistance or type 2 diabetes develops these warning signs, and not everyone who has these symptoms necessarily has type 2 diabetes.

Increased hunger, even after eating  
Unexplained weight loss  
Increased thirst, dry mouth, and frequent urination  
Feeling very tired  
Blurred vision  
Slow healing of sores or cuts  
Dark velvety or ridged patches of skin, especially on the back of the neck or under the arms  
Irregular periods, no periods, and/or excess facial and body hair growth in girls  
High blood pressure or abnormal blood fats levels

## **Type 2 Diabetes Prevention Methods and Treatments**

Healthy lifestyle choices can help prevent and treat type 2 diabetes. Even with a family history of diabetes, eating healthy foods in the correct amounts and exercising regularly can help children achieve or maintain a normal weight and normal blood glucose levels.

**Eat healthy foods.** Make wise food choices. Eat foods low in fat and calories.

**Get more physical activity.** Increase physical activity to at least 60 minutes every day.

**Take medication.** If diet and exercise are not enough to control the disease, it may be necessary to treat type 2 diabetes with medication.

The first step in treating type 2 diabetes is to visit a doctor. A doctor can determine if a child is overweight based on the child's age, weight, and height. A doctor can also request tests of a child's blood glucose to see if the child has diabetes or pre-diabetes (a condition which may lead to type 2 diabetes).

## **Types of Diabetes Screening Tests That Are Available**

**Glycated hemoglobin (A1C) test.** A blood test measures the average blood sugar level over two to three months. An A1C level of 6.5 percent or higher on two separate tests indicates diabetes.

**Random (non-fasting) blood sugar test.** A blood sample is taken at a random time. A random blood sugar level of 200 milligrams per deciliter (mg/dL) or higher suggests diabetes. This test must be confirmed with a fasting blood glucose test.

**Fasting blood sugar test.** A blood sample is taken after an overnight fast. A fasting blood sugar level less than 100 mg/dL is normal. A level of 100 to 125 mg/dL is considered pre-diabetes. A level of 126 mg/dL or higher on two separate tests indicates diabetes.

**Oral glucose tolerance test.** A test measuring the fasting blood sugar level after an overnight fast with periodic testing for the next several hours after drinking a sugary liquid. A reading of more than 200 mg/dL after two hours indicates diabetes.

Type 2 diabetes in children is a preventable/treatable disease and the guidance provided in this information sheet is intended to raise awareness about this disease. Contact your student's school nurse, school administrator, or health care provider if you have questions.